

5 SEM TDC MTMH (CBCS) C 11

2 0 2 2

(Nov/Dec)

MATHEMATICS

(Core)

Paper : C-11

(Multivariate Calculus)

Full Marks : 80

Pass Marks : 32

Time : 3 hours

*The figures in the margin indicate full marks
for the questions*

1. (a) Let partial derivatives of a function of two variables exist. Does it imply that the function is continuous? 1
- (b) Find $\frac{\partial f}{\partial x}$, where $f(x, y) = e^{x^2 + xy}$. 1
- (c) Show that

$$f(x, y) = \begin{cases} \frac{2xy}{x^2 + y^2} & ; (x, y) \neq (0, 0) \\ 0 & ; (x, y) = (0, 0) \end{cases}$$

is continuous at every point, except the origin (0,0). 3

Or

Using definition, show that the function

$$f(x, y) = \begin{cases} \frac{xy}{\sqrt{x^2 + y^2}} & ; (x, y) \neq (0, 0) \\ 0 & ; (x, y) = (0, 0) \end{cases}$$

is continuous at the origin.

(d) Find

$$\frac{\partial^3 u}{\partial z \partial y \partial x} \text{ and } \frac{\partial^3 u}{\partial x^2 \partial y}$$

$$\text{if } u = \frac{x}{y+2z}.$$

2. (a) Write True or False :

"If a function $f(x, y)$ is continuous at (x_0, y_0) , then f is differentiable at (x_0, y_0) ."

(b) Use chain rule to find the derivative of $w = xy$ with respect to t along the path $x = \cos t, y = \sin t$.

(c) Find the values of $\frac{\partial z}{\partial x}$ and $\frac{\partial z}{\partial y}$ at the point (π, π, π) for the function

$$\sin(x + y) + \sin(y + z) + \sin(x + z) = 0$$

P23/435

(Continued)

Or

Find the derivative of $f(x, y, z) = x^3 - xy^2 - z$ at $\rho_0(1, 1, 0)$ in the direction of $\vec{v} = 2\hat{i} - 3\hat{j} + 6\hat{k}$. In what direction does f increase most rapidly at ρ_0 ?

3. (a) Find the plane, tangent to the surface

$$z = x \cos y - ye^x \text{ at } (0, 0, 0).$$

(b) Find the local extreme values of

$$f(x, y) = 3y^2 - 2y^3 - 3x^2 + 6xy$$

(c) Find the points on the hyperbolic cylinder $x^2 - z^2 = 1$ that are closest to the origin.

Find the maximum and minimum values of the function $f(x, y) = 3x + 4y$ on the circle $x^2 + y^2 = 1$.

4. (a) Define gradient vector of $f(x, y)$ at a point.

(b) Show that

$$\vec{f}(x, y, z) = (y^2 z^3)\hat{i} + (2xy z^3)\hat{j} + (3xy^2 z^2)\hat{k}$$

is a conservative vector field.

P23/435

(Turn Over)

(c) Calculate the curl \vec{f} , where

$$\vec{f} = \frac{x\hat{i} + y\hat{j} + z\hat{k}}{\sqrt{x^2 + y^2 + z^2}}$$

2

5. (a) State Fubini's theorem of first form.

1

(b) Evaluate

$$\iint_R f(x, y) dx dy \text{ for } f(x, y) = 1 - 6x^2y^2,$$

$$R : 0 \leq x \leq 1 \text{ and } -2 \leq y \leq 2.$$

2

(c) Prove that

$$\iint_R e^{x^2 + y^2} dy dx = \frac{\pi}{2} (e - 1)$$

where R is the semicircular region bounded by the x -axis and the curve $y = \sqrt{1 - x^2}$.

3

6. (a) Define volume of a region in space.

2

(b) Find $\int_0^2 \int_0^2 \int_0^2 xyz dx dy dz$.

2

(c) Find the volume of the region D enclosed by the surfaces $z = x^2 + 3y^2$ and $z = 8 - x^2 - y^2$.

5

P23/435

(Continued)

Or

Evaluate the following integral by changing the order of the integration in an appropriate way :

$$\int_0^1 \int_0^1 \int_{2y}^2 \frac{4 \cos(x^2)}{2\sqrt{z}} dx dy dz$$

7. (a) Write the formula for triple integral in spherical coordinates.

1

(b) Evaluate :

$$\int_0^\pi \int_0^\pi \int_0^{\sqrt{3-r^2}} dz r dr d\theta$$

4

Or

Find a spherical coordinate equation for the sphere $x^2 + y^2 + (z - 1)^2 = 1$.

8. (a) Define Jacobian of a function of two variables.

1

(b) Evaluate :

3

$$\iint_{x^2 + y^2 \leq a^2} (x^2 + y^2) dx dy$$

P23/435

(Turn Over)

(c) Find the value of

$$\int_C \{(x+y^2)dx + (x^2 - y)dy\}$$

taken in the clockwise sense along the closed curve C formed by $y^3 = x^2$ and the chord joining $(0, 0)$ and $(1, 1)$.

3

Or

Evaluate $\int_C (xy + y + z)ds$ along the curve

$$\vec{r}(t) = 2t\hat{i} + t\hat{j} + (2-2t)\hat{k}, \quad 0 \leq t \leq 1.$$

9. (a) Define line integrals of a vector field. 1

(b) Find the circulation of the field $\vec{F} = (x-y)\hat{i} + x\hat{j}$ around the circle $\vec{r}(t) = (\cos t)\hat{i} + (\sin t)\hat{j}$, $0 \leq t \leq 2\pi$. 3

(c) State and prove the fundamental theorem of line integrals. 4

Or

A fluid's velocity field is $\vec{F} = x\hat{i} + z\hat{j} + y\hat{k}$. Find the flow along the helix $\vec{r}(t) = (\cos t)\hat{i} + (\sin t)\hat{j} + t\hat{k}$, $0 \leq t \leq \frac{\pi}{2}$.

10. (a) Define Green's theorem in Tangential form. 1

(b) Evaluate

$$\oint_C (y^2 dx + x^2 dy)$$

using Green's theorem, where C is the triangle bounded by $x=0$, $x+y=1$, $y=0$.

3

(c) State and prove Stoke's theorem. 6

Or

Evaluate $\int_C \vec{F} \cdot d\vec{r}$ by using Stoke's theorem, if $\vec{F} = x^2\hat{i} + 2xy\hat{j} + z^2\hat{k}$ and C is the ellipse $4x^2 + y^2 = 4$ in the xy plane, counterclockwise when viewed from above.

(d) Use Divergence theorem to find the outward flux of \vec{F} across the boundary of the region D , where

$$\vec{F} = (y-x)\hat{i} + (z-y)\hat{j} + (y-x)\hat{k}$$

and D is the cube bounded by the planes $x = \pm 1$, $y = \pm 1$ and $z = \pm 1$. 5

★★★